



Manufactured By: HYPHEN BioMed

**BIOPHEN Act. PC-r CONTROL PLASMA**  
# A223405

For in vitro diagnostic use only

**BIOPHEN Act. PC-r CONTROL PLASMA**  
# A223405

For in vitro diagnostic use only

Lot : ..... - Exp. : .....

Lot : ..... - Exp. : .....

Control	[C]	Acceptance range
Act PC-r (ratio)*	.....	≤ 1.80
ActPCr (%FVL)	.....	.....-.....

Control	[C]	Acceptance range
Act PC-r (ratio)*	.....	≤ 1.80
ActPCr (%FVL)	.....	.....-.....

\* Note: This is an indicative value that depends on the method and the reagents used. The acceptance range is ≤ 1.80.

Note: For information, clotting times obtained depend on the method and the reagents used, and are of 25 to 40 sec for CT1, and of 45 to 70 sec for CT2, when using Hemoclot FV-L device (ref ACK061K/L) with water bath or STA.

\*Note: This is an indicative value that depends on the method and the reagents used. The acceptance range is ≤ 1.80.

Note: For information, clotting times obtained depend on the method and the reagents used, and are of 25 to 40 sec for CT1, and of 45 to 70 sec for CT2, when using Hemoclot FV-L device (ref ACK061K/L) with water bath or STA.

**BIOPHEN Act. PC-r CONTROL PLASMA**  
# A223405

For in vitro diagnostic use only

**BIOPHEN Act. PC-r CONTROL PLASMA**  
# A223405

For in vitro diagnostic use only

Lot : ..... - Exp. : .....

Lot : ..... - Exp. : .....

Control	[C]	Acceptance range
Act PC-r (ratio)*	.....	≤ 1.80
ActPCr (%FVL)	.....	.....-.....

Control	[C]	Acceptance range
Act PC-r (ratio)*	.....	≤ 1.80
ActPCr (%FVL)	.....	.....-.....

\* Note: This is an indicative value that depends on the method and the reagents used. The acceptance range is ≤ 1.80.

Note: For information, clotting times obtained depend on the method and the reagents used, and are of 25 to 40 sec for CT1, and of 45 to 70 sec for CT2, when using Hemoclot FV-L device (ref ACK061K/L) with water bath or STA.

\* Note: This is an indicative value that depends on the method and the reagents used. The acceptance range is ≤ 1.80.

Note: For information, clotting times obtained depend on the method and the reagents used, and are of 25 to 40 sec for CT1, and of 45 to 70 sec for CT2, when using Hemoclot FV-L device (ref ACK061K/L) with water bath or STA.

Form AH121  
02-2010



7768 Service Center Drive • West Chester OH 45069

Phone: 513.770.1991

Toll Free: 866.783.3797

Fax: 513.573.9241

Email: info@aniara.com

www.aniara.com

**BIOPHEN Act. PC-r CONTROL PLASMA**  
**# A223405**

Abnormal control plasma for Act. PC-r assay quality control

For *in vitro* diagnostic use only

CE

**BIOPHEN Act. PC-r CONTROL PLASMA**  
**# A223405**

Abnormal control plasma for Act. PC-r assay quality control

For *in vitro* diagnostic use only

CE

F.820.09/BI/3405/US /2

F.820.09/BI/3405/US /2

**BIOPHEN Act. PC-r CONTROL PLASMA**  
**# A223405**

Abnormal control plasma for Act. PC-r assay quality control

For *in vitro* diagnostic use only

CE

**BIOPHEN Act. PC-r CONTROL PLASMA**  
**# A223405**

Abnormal control plasma for Act. PC-r assay quality control

For *in vitro* diagnostic use only

CE

F.820.09/BI/3405/US /2

Form AH121  
02-2010

F.820.09/BI/3405/US /2

**ANIARA**

7768 Service Center Drive • West Chester OH 45069

Phone: 513.770.1991

Toll Free: 866.783.3797

Fax: 513.573.9241

Email: [info@aniara.com](mailto:info@aniara.com)

[www.aniara.com](http://www.aniara.com)