

U-CyTech BV

Yalelaan 48
3584 CM Utrecht
The Netherlands
P +31.30.253 5960
F +31.30.253 9344
INFO@ucytech.com
www.ucytech.com

Data sheet Monkey IL-17A ELISA antibody pair; 20-plate format

Cat. No.: CT551-20

Coating antibodies (4 vials)

Product: Monoclonal antibody to monkey interleukin 17A (IL-17A)
Isotype: Mouse IgG_{2b}
Production: *In vitro* using serum free medium
Purification: Ion exchange chromatography
Buffer: Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose
Application: Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50 µl/well).

Detection antibodies (4 vials)

Product: Biotinylated monoclonal antibody to monkey interleukin 17A (IL-17A)
Isotype: Mouse IgG₁
Production: *In vitro* using serum free medium
Purification: Protein A-affinity chromatography
Labeling: With Biotin-7-NHS (N-hydroxysuccinimide)
Buffer: Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).

Standards (10 vials)

Product: Recombinant cynomolgus monkey interleukin 17A (IL-17A)
Application: Cytokine standard in an ELISA system
Reconstitution: Dissolve the contents of one vial by injection of 0.5 ml distilled water into the vial. Use immediately.

Conjugate (4 vials)

Product: SPP conjugate (Streptavidin-HRP)
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).
The product should be used in combination with TMB substrate.

General

Sensitivity: 1 pg/ml
Specificity: Validated for detecting natural rhesus macaque and baboon IL-17A. Standards are calibrated with natural rhesus macaque IL-17A.
Sterility: Membrane filtered (0.2 µm)
Stability: Lyophilized SPP conjugate is stable for at least one year at -20°C in the dark, the other lyophilized products are stable for more than one year at 4°C. After reconstitution, the antibodies are stable for one year at 4°C (if kept sterile) and SPP for minimal one year at -20°C in the dark. The reconstituted standard preparation should be used immediately.

PROUDLY DISTRIBUTED BY

ANIARA
ANIARA DIAGNOSTICA LLC

+1 (513) 770-1991 | +1 (866) 783-3797
7768 Service Center Drive
West Chester, OH 45069
info@aniara.com

For research use only

Version 160629