

Data sheet **Marmoset IFN- γ ELISA antibody pair; 20-plate format**

Cat. No.: CT770-20

Coating antibodies (4 vials)

Product: Monoclonal antibody to marmoset interferon gamma (IFN- γ)
Isotype: Mouse IgG₁
Production: *In vitro* using serum free medium
Purification: DEAE ion exchange chromatography
Buffer: Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose
Application: Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50 μ l/well).

Detection antibodies (4 vials)

Product: Biotinylated monoclonal antibody to marmoset interferon gamma (IFN- γ)
Isotype: Mouse IgG₁
Production: *In vitro* using serum free medium
Purification: DEAE ion exchange chromatography
Labeling: With Biotin-7-NHS (N-hydroxysuccinimide)
Buffer: Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 μ l/well).

Standards (10 vials)

Product: Recombinant marmoset interferon gamma (IFN- γ)
Application: Cytokine standard in an ELISA system
Reconstitution: Inject 0.5 ml distilled water into the vial. Use immediately.

Conjugate (4 vials)

Product: SPP conjugate (Streptavidin-HRP)
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 μ l/well).
The product should be used in combination with TMB substrate.

General

Sensitivity: 1 pg/ml
Specificity: Validated for detecting natural and recombinant marmoset IFN- γ . Standards are calibrated with recombinant marmoset IFN- γ .
Sterility: Membrane filtered (0.2 μ m)
Stability: Lyophilized SPP conjugate is stable for at least one year at -20°C in the dark, the other lyophilized products are stable for more than one year at 4°C. After reconstitution, the antibodies are stable for one year at 4°C (if kept sterile) and SPP for minimal one year at -20°C in the dark. The reconstituted standard preparation should be used immediately.

PROUDLY DISTRIBUTED BY

ANIARA
ANIARA DIAGNOSTICA LLC

+1 (513) 770-1991 | +1 (866) 783-3797
7768 Service Center Drive
West Chester, OH 45069
info@aniara.com